



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

# Journal Pre-proof

Seroconversion rate after vaccination against COVID-19 in cancer patients - a systematic review

C. Corti, G. Antonarelli, F. Scotté, J.P. Spano, J. Barrière, J.M. Michot, F. André, G. Curigliano

PII: S0923-7534(21)04550-6

DOI: <https://doi.org/10.1016/j.annonc.2021.10.014>


Reference: ANNONC 782

To appear in: *Annals of Oncology*

Received Date: 2 September 2021

Revised Date: 3 October 2021

Accepted Date: 23 October 2021



Please cite this article as: Corti C, Antonarelli G, Scotté F, Spano J, Barrière J, Michot J, André F, Curigliano G, Seroconversion rate after vaccination against COVID-19 in cancer patients - a systematic review, *Annals of Oncology* (2021), doi: <https://doi.org/10.1016/j.annonc.2021.10.014>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 European Society for Medical Oncology. Published by Elsevier Ltd. All rights reserved.

**Title:** Seroconversion rate after vaccination against COVID-19 in cancer patients - a systematic review

**Author names:** C. Corti<sup>1,2</sup>, G. Antonarelli<sup>1,2</sup>, F. Scotté<sup>3,4</sup>, JP. Spano<sup>5</sup>, J. Barrière<sup>6</sup>, JM. Michot<sup>7</sup>, F. André<sup>8</sup>, G. Curigliano<sup>1,2</sup>

**Author affiliations:**

<sup>1</sup> Division of New Drugs and Early Drug Development for Innovative Therapies, European Institute of Oncology, IRCCS, Milan, Italy

<sup>2</sup> Department of Oncology and Haematology (DIPO), University of Milan, Milan, Italy

<sup>3</sup> Gustave Roussy Cancer Campus, Villejuif, France

<sup>4</sup> Département Interdisciplinaire d'Organisation des Parcours Patients, Gustave Roussy, Villejuif, France.

<sup>5</sup> APHP-Sorbonne Université, Institut Pierre Louis d'Epidemiologie et de Santé Publique INSERM 1136, Paris, France

<sup>6</sup> Department of Medical Oncology, Clinique Saint-Jean, Cagnes-sur-Mer, France

<sup>7</sup> Drug Development Department (DITEP) Gustave Roussy Cancer Campus, Villejuif, France

<sup>8</sup> Predictive biomarkers and novel therapeutic strategies Group, Institut Gustave Roussy, University of Paris Sud, INSERM 981, Université Paris Saclay, Villejuif, France.

**Corresponding author:** Prof. Giuseppe Curigliano, MD, PhD, Division of New Drugs and Early Drug Development for Innovative Therapies, European Institute of Oncology, IRCCS, Via Ripamonti 435, 20141, Milan, Italy. Phone number: +39 0257489599. (E-mail: giuseppe.curigliano@ieo.it).

30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57

**Authors' ORCIDs:**

Chiara Corti: 0000-0002-0359-0267  
Gabriele Antonarelli: 0000-0002-9758-3665  
Florian Scotté: 0000-0001-6175-7208  
Jean-Philippe Spano: 0000-0002-3921-2479  
Jérôme Barrière: 0000-0003-0961-8002  
Jean-Marie Michot: 0000-0003-3482-3331  
Fabrice André: 0000-0001-5795-8357  
Giuseppe Curigliano: 0000-0003-1781-2518

**Words count: 197/300 (abstract) + 5224 (review).**

58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83

**ABSTRACT (197/300 words):**

*Background.* Coronavirus disease 2019 (COVID-19) has affected more than 210 million people worldwide. An optimal therapeutic approach for COVID-19 remains uncertain, to date. Since the history of cancer was linked to higher mortality rates due to COVID-19, the establishment of a safe and effective vaccine coverage is crucial in these patients. However, patients with cancer were mostly excluded from vaccine candidates' clinical trials. This systematic review aims to investigate the current available evidence about the immunogenicity of COVID-19 vaccines in patients with cancer (PsC).

*Patients and methods.* All prospective studies that evaluated safety and efficacy of vaccines against severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) were included, with immunogenicity after the first and the second dose as the primary endpoint, when available.

*Results.* Vaccination against COVID-19 for PsC seems overall safe and immunogenic after well-conducted vaccinations schedules. Yet, the seroconversion rate remains lower, lagged or both compared to the general population. Patients with hematologic malignancies, especially those receiving B cell depleting agents in the last 12 months are the most at risk of poor seroconversion.

*Conclusion.* A tailored approach to vaccination may be proposed to PsC, especially on the basis of the type of malignancy and of the specific oncologic treatments received.

**KEYWORDS:** COVID19, Sars-CoV-2, vaccine, immunogenicity, cancer, seroconversion.

84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108

**HIGHLIGHTS:**

1. History of cancer is linked to higher mortality rates due to COVID-19. **(69/125)**
2. Data on the immunogenicity of COVID-19 vaccines in patients with cancer (PsC) are scarce. **(88/125)**
3. This review included prospective studies that evaluated immunogenicity of vaccines against COVID-19 in PsC. **(106/125)**
4. Vaccination seems overall safe and immunogenic after well-conducted vaccinations schedules. **(90/125)**
5. Hematologic patients receiving B cell depleting agents in the last 12 months are the most at risk of poor seroconversion. **(121/125)**

109

110 **MANUSCRIPT** (5224 words)111 **INTRODUCTION**

112 Since the first reports of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection,  
113 coronavirus disease 2019 (COVID-19) has affected more than 210 million people worldwide (1).  
114 Besides oxygen therapy and positive pressure ventilation, glucocorticoids, especially  
115 dexamethasone, showed a mortality benefit in patients requiring respiratory support (2). Despite  
116 different therapeutic approaches being investigated, an optimal treatment for COVID-19 remains  
117 uncertain (**Figure 1**) (3-5).

118 Nevertheless, global efforts have established an effective vaccine strategy and, because history of  
119 cancer is linked to higher mortality rates due to COVID-19, an effective vaccine coverage is crucial  
120 in this population (6-13). However, clinical trials investigating COVID-19 vaccine candidates mostly  
121 excluded patients with cancer (PsC). So, international COVID-19 vaccination guidelines for this  
122 population were initially based on expert opinions, on studies designed to test other vaccines and  
123 on initial real-world data reports (14, 15).

124 This systematic review aims to investigate the current available evidence about immunogenicity of  
125 COVID-19 vaccines currently administered to PsC.

126

127 **METHODS**

128 A systematic review of the literature was performed on August 16th, 2021. The relevant studies were  
129 searched through Medline (via PubMed) and Embase, with no language or time restriction. The  
130 databases were searched (CC) using the mapped terms ["cancer" OR "tumor" OR "malignancy"]  
131 AND "vaccine" AND ["COVID" or "SARS-CoV-2"] and the exploded MeSH terms "COVID-19  
132 Vaccines". Two reviewers double-screened independently titles and abstracts (CC, GA). A third  
133 author functioned as tiebreaker, in case of disagreements (GC). The reference lists of the most  
134 relevant papers were selected for snowballing (CC, GA). The Preferred Reporting Items for  
135 Systematic Reviews and Meta-Analyses (PRISMA) methodology was applied, to depict the flow of  
136 studies through each phase (ph) of the review process.

We assessed the quality of included studies using the Quality Assessment Tool for Observational Cohort and Cross-Sectional Studies (NIH). Results were rated as FAIR, when a total of 5-10 points were assigned the study and as GOOD if a total of  $\geq 11$  points were assigned on the basis of 14 quality-assessment queries. A comprehensive summary and specifics of the quality assessment are provided in **Supplementary Table S1**.

Data extraction was performed by one reviewer (CC) and independently checked by other two authors (GA, GC). We included all prospective studies, evaluating as primary endpoint the immunogenicity of vaccines against SARS-CoV-2 in PsC. Findings which did not fulfil the above-mentioned criteria were excluded. Safety was investigated as secondary endpoint, as the incidence of adverse events (AEs), if reported in the clinical study.

We retrieved supplementary information about study design, population size and cancer types (if available). When a single study had resulted in multiple publications, we prioritized the most updated report, unless the reported endpoint was not relevant.

Substantial heterogeneity of study designs and outcome measures did not allow to perform a meta-analysis; therefore, a narrative synthesis was conducted without performing additional statistical- or sensitivity analyses by a specific software or without additional feasibility assessment.

## RESULTS

The systematic research of the literature returned 2526 records. After checking for duplicates, a total of 1607 records were obtained. After critical appraisal, a final amount of 36 studies met the above-mentioned criteria, as depicted in **Figure 2**. The investigations have been performed in the World Health Organization (WHO) Region of the Americas ( $n = 7$ ) and in the WHO European Region (i.e., United Kingdom,  $n = 8$ ; Germany,  $n = 1$ ; Denmark,  $n = 1$ ; Italy,  $n = 3$ ; France,  $n = 3$ ; Greece,  $n = 3$ ; Switzerland,  $n = 1$ ; Israel,  $n = 6$ ; Turkey,  $n = 1$ ; Lithuania,  $n = 1$ ; Netherlands,  $n = 1$ ). The median and the mean number of PsC included in each study were 114.5 (min: 16; max: 1503) and 257.2, respectively, overall accounting for 9260 patients. As for vaccines platforms, Pfizer-BioNTech (BNT) and Moderna (MDN) vaccines were administered in 33 (91.6%) and in 13 (36.1%) clinical studies, respectively. Viral vector-based vaccines include Oxford/AstraZeneca (OxA) and Janssen vaccines.



165 The former was administered in 9 clinical studies (25%), whereas the latter was evaluated only in  
166 one study. The study conducted in Turkey administered CoronaVac, an inactivated COVID-19  
167 vaccine (16).

168 Across all the studies, immunogenicity of COVID-19 vaccines was defined as the proportion of PsC  
169 who seroconverted to Spike proteins. Median anti-Spike antibody (Ab) titers, detection of neutralizing  
170 antibodies (NAbs) and cellular immune responses were investigated as secondary or exploratory  
171 endpoints.

172 Eight studies reported data only after the first dose; twenty studies reported data only after the  
173 second dose; nine studies reported data after both the first and the second dose. The seroconversion  
174 rate ranged widely after the first dose, i.e., from 11% to 87.5%, overall; from 11% to 87.5% for  
175 hematologic patients; from 25% to 67% for patients with solid tumors. However, seroconversion data  
176 were collected at non-uniform time points after the first dose, across all the studies (median: 3 weeks;  
177 min: 1 week; max: 5 weeks).

178 As for the second dose, the seroconversion rate ranged widely as well, from 7.3% to 100%.  
179 Specifically, it ranged from 7.3% to 88.8%, for hematologic patients and from 47.5% to 100% for  
180 patients with solid tumors. Similarly, seroconversion data were collected at non-uniform time points  
181 after the second dose, across all the studies (median: 3 weeks; min: 1 week; max: 16 weeks).

182 When safety measures were described, the incidence of AEs was the most commonly reported  
183 outcome, even though 22 studies (61.1%) did not extensively report safety information. Conversely,  
184 2 studies (5.56%) reported the incidence of AEs only after the first dose; five studies (13.9%) reported  
185 AEs only after the second dose. Finally, in 7 studies (20.6%) AEs were reported after both the first  
186 and the second dose. Overall, COVID-19 vaccines resulted safe and well tolerated, with no vaccine-  
187 related deaths. Any-grade AEs ranged between 9.7% and 87% after the first dose and between 23%  
188 and 85% after the second dose. The most commonly reported any-grade AEs were local pain (range:  
189 7.4% - 69%, I dose; range: 32.3% - 67.2%, II dose) and fatigue (range: 4.2% - 47.6%, I dose; range:  
190 3% - 23.4, II dose).

191 Patients with solid tumors were included in 15 studies (41.7%), whereas hematologic malignancies  
192 were represented in 28 studies (77.8%). Twenty-one studies (58.3%) were exclusively focused on

hematologic patients. Six studies specifically focused on PsC on active treatments, namely cytotoxic agents, B-cell depleting agents, Janus kinase inhibitors (JAKi) and immune checkpoint inhibitors (ICIs) (17-22). A comprehensive summary of the studies included is provided in **Supplementary Table S2**.

### Cancer Type

Available evidence suggests that vaccines, besides being generally safe and well tolerated, may have a compromised activity, especially in the case of hematologic malignancies (**Figure 3**) (24). In this regard, a prospective observational study included 151 PsC (95 with a solid tumor and 56 with a hematologic malignancy), and 54 healthy controls (**Supplementary Table S2**) (25). In an interim analysis, the proportion of patients with positive anti-Spike IgG titers at approximately 21 days following the first dose was 94% for the healthy controls, compared with 38% of those with solid tumors, and 18% of those with hematologic malignancies (25). Considering patients with available blood samples two weeks after the second dose, 95% of the patients with solid tumors and 60% of those with hematologic malignancies showed seropositivity, in comparison with 100% of healthy controls. Another study evaluated 200 patients, of which 134 harbored solid tumors and the remaining 66 had a hematologic diagnosis. Vaccination was carried out with MDN (62/200), BNT (115/200) and Janssen (20/200). Although the overall seroconversion rate reached 94%, hematologic malignancies revealed a significantly lower rate (85%), particularly among those receiving B-cell depleting therapies and following hematopoietic cell transplantation (HCT) (73%) (21). For example, a detailed study highlighted that anti-CD20 Abs, Bruton tyrosine kinase inhibitors (BTKi), JAKi and B-cell lymphoma 2 (bcl-2) inhibitors seemed to electively impact on the Ab response to vaccination (24). Importantly, when vaccination was administered 12 months after the last treatment, serological responses improved (24). Consistently, initial findings from the CAPTURE study, a prospective longitudinal cohort study of SARS-CoV-2 infection and COVID-19 vaccine-induced immunity, were recently presented (26). Seroconversion rates for anti-Spike (S1) Abs following 2 doses were 85% and 54% for patients with solid tumors and hematologic malignancies, respectively. This study specifically focused on neutralizing antibodies (NAbs), describing lower

detection rates and NAb titers in patients with hematologic malignancies in comparison with patients harboring solid tumors (26). Notably, after natural infection, neutralizing antibodies remained stable, unlike anti-Spike (S1) Abs that waned over time (26). Similar Ab production in PsC was shown in other studies (**Supplementary Table S2**) (19, 24, 27, 28).

#### *Multiple myeloma (MM)*

Immunogenicity of COVID-19 vaccines has been investigated also in patients with specific hematologic conditions, such as MM. Among 103 patients (96 with active MM) who received mRNA-based vaccines, only 45% of active MM patients developed an adequate immune response, while 22% had a partial response, when stratified according to Ab titer (29). Conversely, smoldering MM patients ( $n = 7$ ) responded better (29). Lower anti-Spike Abs levels were associated with older age, impaired renal function, low lymphocyte counts, reduced uninvolved immunoglobulin levels, > second line of treatment, and absence of complete remission (29). The increased risk of poor seroconversion has been highlighted also in another study, though only focusing on the first vaccine dose (30). Other studies focusing on plasma cell disorders reported similar results (31, 32). Consistently, two retrospective analyses investigated seroconversion in response to COVID-19 vaccines in a series of 320 and 23 fully immunized MM patients, respectively (33, 34). In one study, individuals were assessed for serologic response at least 10 days after receiving the second dose of a mRNA-based vaccine. Although 84% of patients mounted a measurable Ab response, the serologic titer varied by three orders of magnitude (range: 5 - 7882 AU/mL, median: 149 AU/mL) (33). Similarly, in the second study, the seroconversion rate reached 74% with a median anti-Spike titer or 4.9 UI/mL (range: 0-1028) (34).

#### *Myelodysplastic (MDS) and myeloproliferative (MPN) neoplasms*

MPNs are associated with a pro-inflammatory state and dysregulation of pivotal natural killer cell (NK), regulatory T cell (Tregs) and effector T cell function (20). Two prospective studies evaluated the immune response after the first dose of COVID-19 vaccine in patients with MPNs. In one study only BNT was administered (20). In the second study, both mRNA-based and viral vector-based

249 vaccines were administered (35). After the first dose, patients with a diagnosis of myelofibrosis (MF)  
 250 ( $n = 9$ ) had significantly higher post-vaccine anti-Spike IgG half maximal effective concentration  
 251 (EC50) as well as neutralizing Ab inhibitory dose (ID)50 titers, compared to patients with other MPN  
 252 subtypes (20). Seroconversion measured >14 days after a single dose was only 58%, that is  
 253 significantly lower than the one observed in health-care professionals (HCPs) of similar age (97%).  
 254 The median anti-spike Ab titer was also significantly lower in MPN/MDS patients (i.e., 630 vs 75  
 255 AU/ml,  $p < 0.0001$ ) (35). When focusing on disease subgroups, the seroconversion rate was highest  
 256 in patients with chronic myeloid leukemia (CML, 75%), with no difference according to which vaccine  
 257 was administered (35). Another study evaluated seroconversion rates at 5 weeks after the first BNT  
 258 dose, thus also including patients receiving the second dose (36). Seroprotection rate at cut-off of  
 259 15 AU/mL was 100% in controls compared to 88% in MPN patients ( $p = 0.038$ ) (36).

260

### 261 *Lymphoma*

262 Patients diagnosed with lymphoma are at particular high risk of severe COVID-19 (15). Recently,  
 263 one prospective observational study evaluated the humoral immune response to BNT in a cohort of  
 264 148 patients harboring B-cell non-Hodgkin lymphoma (B-NHL). Of those, 47% displayed an  
 265 aggressive disease, whereas 53% had an indolent malignancy (37). Of note, 37% of patients were  
 266 receiving active treatment. Ab titer was measured 2-3 weeks after the second vaccine dose.  
 267 Seroconversion was achieved in 49% of B-NHL patients versus a 98.5% rate achieved in healthy  
 268 controls ( $p < 0.001$ ) (37).

269 In the interim analysis of the PROSECO study participants received either OxA or BNT, with two  
 270 doses given 10-12 weeks apart. A total of 129 patients were enrolled. Of those, 12 patients (9%) had  
 271 Hodgkin lymphoma (HL), 34 (26%) had aggressive B-NHL, 79 (61%) had indolent B-NHL and 4 (3%)  
 272 had peripheral NK/T cell lymphoma (38). Notably, 52 (44%) of 119 participants with lymphoma were  
 273 on active treatment (38). Twenty-two (72%) of 31 participants after one dose of vaccine and 20/33  
 274 (61%) participants after two doses did not produce detectable anti-spike IgG Abs. Among the  
 275 lymphoma patients who were not on active treatment, 6/6 (100%) patients with HL and 13/16 (81%)  
 276 with aggressive B-NHL developed an immune response comparable to that of healthy individuals

(38). Thirty-two (89%) of 36 participants with an indolent B-NHL who were not on active treatment showed detectable Abs after two vaccine doses. However, their Ab titer was reduced in comparison with the levels observed in participants with HL and aggressive B-NHL that were either treatment-naïve or with completion of treatment > 3 years prior to vaccination.

#### *Chronic lymphocytic leukemia (CLL)*

Compared with other hematologic malignancies, the Ab response appears particularly impaired in CLL patients. A prospective study that compared serologic response with BNT between matched cohorts of 52 patients and 52 healthy subjects showed that CLL patients had a lower serologic response rate (52% versus 100%) than healthy controls ( $p < 0.001$ ) (39). When focusing on the entire cohort of 167 CLL patients, the Ab response rate was only 39.5%, with younger age, lack of active treatment and early disease stage associated with better seroconversion rates. Other studies focusing on CLL reported similar results, suggesting that humoral response may be particularly affected by disease activity itself (34, 40, 41).

#### *Solid tumors*

A prospective study investigated the serologic status of the BNT in a cohort of patients with solid tumors on active treatment ( $n = 232$ ), compared with age-matched HCPs ( $n = 261$ ). In the patient group, 86/232 individuals were tested after the first vaccination dose and 218/232 were tested after the second dose. After the first dose, 25/86 (29%) patients were seropositive compared with 220/261 (84%) healthy controls ( $p < 0.001$ ). After the second dose, the seropositive rate reached 86% (187/218) among the PsC (27). At the latest time point (4 weeks after the second dose), 14% of PsC were seronegative. Specifically, patients with breast cancer (BC) accounted for 29% of the seronegative group and 74% of these individuals were treated with diverse regimens of chemotherapy (CT) (27). However, although specific CT agents may not be directly linked to impaired immunogenicity, the lymphosuppressive potential of some CT regimens may limit seroconversion (27).

Another study including 95 patients with solid cancer and 66 healthy controls reported a seroconversion rate of 87% and 100% in patients and controls, respectively, after a median of 123 days from the second vaccination. However, a significantly lower median titer levels in PsC was found, in comparison with the control group (417 AU/mL versus 1220 AU/mL,  $p < 0.001$ ) (42). In an exploratory multivariate analysis, the co-administration of CT plus immunotherapy (IO) or of IO plus a biological agent resulted the only variable associated with lower IgG titers. A recent pooled analysis including 223 PsC with solid tumors highlighted a higher seroconversion rate (94%), with significantly lower anti-Spike Abs, compared to healthy controls, irrespective of the assay used (19, 28, 43).

### **Type of treatment**

A major unanswered question for PsC is whether vaccine immunogenicity is impacted by the concomitant use of specific drugs. Initial findings from the VOICE trial, focusing on solid tumors, have been recently presented (44). Among patients receiving IO, CT and CT-IO, anti-Spike (S1) IgG seroconversion rates were 99.3%, 97.4% and 100%, respectively. As the authors established a cut-off of 300 BAU/mL for adequate Ab response, seroconversion rates after two vaccine doses dropped to 93.1%, 83.8% and 88.8%, for patients receiving IO, CT and CT-IO, respectively (44). Thus, a significant minority of patients does not develop an adequate Ab response (6.9%, IO; 16.2%, CT; 11.2%, CT-IO). Few other studies focused on individual agents to fully elucidate any potential interaction with the ability to mount a protective immune response. Emerging data are clarifying that, in general, the protective role of two vaccine doses for PsC on certain active treatments may be suboptimal (**Figure 3**) (16, 19, 35, 38, 45, 46).

### ***Endocrine treatment (ET)***

In a recent study, post-vaccination seroconversion rates in patients receiving ET ( $n = 47$ ) resulted high in comparison with other active treatments, reaching a 100% seropositivity rate ( $p = 0.04$ ) (21). Therefore, no major preventative measures or time windows should be implemented in current vaccination campaigns.

### *Cytotoxic chemotherapy*

Increasingly consistent data suggest that among patients receiving systemic CT for solid tumors, there is a high proportion of weakly responsive and unresponsive patients after a single vaccine dose (19). In this regard, immune responses to BNT were evaluated in 52 solid tumor patients on active cytotoxic CT and compared to a control group of 50 healthy individuals (47). Neutralizing Abs were detected in 67% and in 80% of PsC after the first and the second dose, respectively (47). Similar trends were observed as for Abs against the receptor-binding domain (RBD) and the S2 regions of the Spike protein, although they were found to be reduced in comparison to healthy controls (47). Other studies showed similar results, consistent with a link between the lymphosuppressive potential of some CT regimens and a delay or reduction in an effective seroconversion (25, 27, 42, 43, 48).

### *Targeted therapies*

Evidence addressing the seroconversion rates in PsC on specific targeted agents is lacking (19, 49). Compounds causing lymphopenia or specific B-cell-depleting agents are emerging as major responsible for an impaired protection from vaccines (24). However, the long-term immunologic effects of B-cell depletion and the characteristics of B cell reconstitution, especially in lymphoid malignancies, are not well defined, despite the widespread usage of B cell directed therapies (46).

*B-cell depleting therapies.* Treatment with B cell directed agents - e.g., rituximab and obinutuzumab (anti-CD20), ibrutinib (BTKi) - may negatively impact the production of Abs in response to COVID-19 vaccines, especially in lymphoid malignancies, due to B-cell depletion and/or disruption of the B-cell receptor signaling pathway (46). In addition, the recovery of the memory B cell pool has been shown to be delayed in lymphomas, remaining below normal controls at 1 year post rituximab (46, 50).

In a cohort of 149 B-NHL patients, 37% were actively treated with a rituximab/obinutuzumab (R/Obi)-based regimen for either induction or maintenance, whereas 44% had last been treated with R/Obi 6 months before COVID-19 vaccination. Seroconversion was achieved in 25/28 (89%) treatment-naïve patients, in 4/55 (7.3%) R/Obi patients and in 43/65 (66.7%) patients receiving the last dose



360 of the B-cell depleting regimen 6 months prior to vaccination. Multivariate analysis revealed that a  
361 longer time window since last R/Obi exposure and an absolute lymphocyte count  $\geq 0.9 \times 10^3/\text{mL}$   
362 predicted seroconversion (37).

363 An optimal window for vaccination in this patient population should be considered. In fact, in another  
364 study focused on lymphomas, seroconversion rates differed between patients who received the last  
365 infusion of an anti-B cell agent within 9 months prior to the COVID-19 vaccine (6/52, 11%) compared  
366 to those who received the last infusion of a B cell depleting agent > 9 months prior to the vaccination  
367 (22/25, 88%) (46). Consistently, in a cohort of CLL patients, none of the 22 individuals who received  
368 therapeutic anti-CD20 Abs in the 12 months prior to COVID-19 vaccination developed neutralizing  
369 Abs after two mRNA-based vaccination, in comparison with 25/55 (46%) patients exposed to anti-  
370 CD20 therapy  $\geq 12$  months prior to vaccination (39). Similar results came from other recent reports,  
371 with a significant 14.2-fold increased risk of non-responding to COVID-19 vaccination (24, 51, 52).  
372 Finally, serologic responses are also impaired in certain populations with MM receiving therapies  
373 against CD38 (e.g., daratumumab, isatuximab) and B cell maturation antigen (anti-BCMA agents,  
374 e.g., belantamab mafodotin and idecabtagene vicleucel) (30, 31, 33, 36).

375 *Janus kinase inhibitors.* JAKi, such as ruxolitinib (RUXO), are currently approved for the treatment  
376 of MF and hydroxyurea resistant/refractory polycythemia vera (17). RUXO is thought to have  
377 profound effects on different cell compartment of the immune system, including T cells, NK cells and  
378 dendritic cells. By inhibiting JAK-signal transducer and activator of transcription (STAT) signaling, a  
379 potential role in reducing inflammatory cytokine production is considered (17). Such features could  
380 explain the increased rate of infection in MPN patients receiving RUXO (17).

381 A prospective study recently assessed the serologic response after the first COVID-19 mRNA-based  
382 vaccine injection in 30 consecutive MPN patients receiving RUXO at a median dose of 20 mg daily.  
383 The Ab response after a first vaccination dose was significantly lower in RUXO-treated patients  
384 compared to healthy controls and to patients not receiving RUXO. All 14 healthy controls were  
385 vaccine-responder (100%), whereas only 33.3% of RUXO-treated patients seroconverted to Spike  
386 protein ( $p < 0.001$ ). As for patients not receiving RUXO, the seroconversion rate was 91.6% (17).  
387 Further studies, hopefully with larger sample sizes, are needed to address whether such



unresponsive status persist after the second vaccine dose, as suggested by the 42% seropositivity rate described in MPN patients using JAKi, after completing the full mRNA-based vaccination schedule (53).

*Cyclin-dependent kinases (CDK) 4/6 inhibitors.* The involvement of the CDK4/6 pathway in immune activation is well-known (54). In cohort of 23 BC patients receiving CDK4/6 inhibitors, neutralizing Ab titers in response to the first dose of vaccine were similar to healthy controls (55). However, anti-Spike Ab titers were low in another study assessing the response after full vaccination, even if the subset of patients was too small ( $n = 5$ ) to draw solid conclusions (21).

*Tyrosine kinase inhibitors (TKI).* To date, there is no evidence that any of the TKIs currently approved in clinical practice can interfere with effective immune responses against SARS-CoV-2 (21). In spite of a small sample size, a recent report on CML patients receiving imatinib showed conserved seroconversion rates (5/6, 83%) (35). Among patients who received other TKIs, namely nilotinib, bosutinib and dasatinib, the seroconversion rate was 66.7% (4/6) (35).

401

#### 402 *Immunotherapy*

*Chimeric antigen receptor (CAR)-T cell therapy.* Among patients with aggressive B-NHL receiving CAR-T cell therapy, 100% (3/3) had no detectable Abs after the first vaccination dose. Only one of these patients developed Abs after the second dose, even if the other two had yet to be tested at the time of the study report. These results were observed although these patients had completed CAR-T cell treatment 11-23 months before vaccination (38).

*Immune checkpoint inhibitors.* An unresolved question for PsC is whether vaccine safety and/or immunogenicity are impacted by ICIs, which stimulate immune system function (22, 56). As for safety, in a study enrolling 134 PsC receiving ICIs, either as monotherapy or in combination with CT, the AEs of COVID-19 vaccination seemed comparable with those of healthy controls. Of note, only the incidence of muscle pain was higher (57). However, there was no immune-related myositis, and COVID-19 vaccination did not appear to exacerbate or cause new immune-related AEs (57).

Concerning immunogenicity, a recent clinical report documented that 15/59 (25%) versus 186/283 (65.7%) PsC developed neutralizing Ab titers after the first dose ( $p < 0.001$ ) (18). Conversely, in a

second report, seroconversion reached 97% (21). Accordingly, the VOICE trial described seroconversion rates of 99.3% and 100% among patients receiving IO and CT-IO, respectively (44). However, a significant minority of patients did not develop an adequate Ab response (6.9%, IO; 16.2%, CT; 11.2%, CT-IO, with an established cut-off of 300 BAU/mL) (44). As an exploratory finding, the CAPTURE trial highlighted a negative impact of ICIs on cellular immune responses, for which further research is warranted (26).

## DISCUSSION

In the general population, the adaptive immune response to SARS-CoV-2 comprises B cells that produce different classes of Abs in order to neutralize the virus, as well as T cells that support Ab production while also directly killing virus-infected cells (15). Although memory B and T cells have been described both in individuals with a natural infection and in vaccinated populations, their specific roles in achieving protective immunity have to be defined yet (49, 58, 59). However, T cells are thought to play an important role in reducing COVID-19 severity (60, 61). Several observations suggest that early SARS-CoV-2 T cell responses are associated with milder COVID-19 (62, 63). In this regard, data from phase III clinical trials investigating COVID-19 vaccines suggest that protection may require low levels of neutralizing Abs and might involve other immune effector mechanisms, including non-neutralizing Abs, T cells and innate immunity (58). Circulating Ab titers did not result to be predictive of T cell memory (59, 64, 65). Additionally, although real world data indicates that vaccine protection against SARS-CoV-2 infection wanes over time, protection against hospitalization and severe disease appears to be preserved (66-68).

For PsC, dissecting the complexity of a protective immune response against SARS-CoV-2 is challenging, considering both the biological differences among cancer types as well as the different treatments received (69). Moreover, since PsC were largely excluded from phase III clinical trials testing vaccine candidates, evidence about protective immune responses came from highly heterogeneous single-center observational studies (**Supplementary Table S2**) (14).

442 Although emerging evidence suggests that simple serological tests for SARS-CoV-2 Abs may not  
443 reflect the complexity and durability of protective immunity against COVID-19, the primary endpoint  
444 of all the studies included was seroconversion to the anti-Spike protein (58, 65, 70).

445 When cellular immunity was investigated, it was considered as an exploratory endpoint (26, 44).  
446 Overall, T cell immune responses seem generally maintained, although reduced, especially in  
447 patients with hematologic malignancies, in comparison to patients with solid tumors and to healthy  
448 individuals, both after COVID-19 infection and after a complete vaccination, with predominance of  
449 CD4+ responses over CD8+ (21, 26, 48, 52, 59). Systemic therapies did not have major effect on  
450 cellular responses, except for higher suppression rates of CD4+ activity among patients treated with  
451 ICIs (26). In the VOICE clinical trial, almost half of the vaccine non-responders and suboptimal  
452 responders to humoral immunity developed a Spike specific T cell response (44). Interestingly, in  
453 the preliminary report of the SOAP-2 vaccine study, the T cell response appeared to be greater than  
454 the B cell immune response after the first dose, but still lower compared to the control group  
455 (**Supplementary Table S2**) (25, 49). Currently, data regarding the clinical efficacy of SARS-CoV-2  
456 vaccines, defined as the incidence of symptomatic or severe COVID-19 in PsC, after full completion  
457 of the vaccination schedule, is still lacking (26). A longer follow up is needed in order to address this  
458 crucial question.

459 With all the described limitations regarding the current understanding of immune responses to  
460 COVID-19 vaccines, the decreased seroconversion rates in PsC, especially in those on active  
461 treatment with B cell depleting agents, could not be neglected (34, 49, 72). Moreover, even if T cell  
462 immunity could be thought to compensate for impaired humoral immunity, impaired T cell responses  
463 in the event of COVID-19 have been observed in some patients under cancer treatment (26, 44, 70,  
464 73). For these reasons, different strategies to enhance vaccine-induced immunity have been  
465 proposed, such as heterologous prime-boost vaccination, a double-dose strategy and a third dose  
466 (49, 74). In the first case, the non-inferiority design of the trial as well as the lack of PsC under active  
467 treatment included, did not allow to draw conclusions about the feasibility of this approach (74). The  
468 double dose strategy is based on literature data and current vaccination practices, particularly in  
469 immunocompromised patients vaccinated against hepatitis B and influenza viruses, although

prospective randomized trials are needed (49, 75-81). Finally, the role of a third dose is being investigated, especially in immunocompromised patients and the elder population (82).

To date, the largest study investigating the clinical efficacy of a booster shot of vaccine was conducted in Israel and included people over 60 years of age (83). The rates of confirmed COVID-19 and severe illness were substantially lower among those who received the booster at least 5 months after the last dose (83). A number of other reports confirmed a benefit on seroconversion for a third mRNA-based vaccine dose, although longer follow-up and evaluation of cellular immune responses are needed to better characterize the impact of additional vaccine doses on the clinical outcomes in patients with impaired immune system (e.g., OCTAVE DUO trial) (52, 72, 80, 82, 84-88). Furthermore, in the CAPTURE clinical trial, previous SARS-CoV-2 infection boosted vaccine-induced responses, lending further support for a third dose in vulnerable populations (26).

In line with these trends, many countries and institutions have already recommended that severely immunocompromised patients (e.g., transplant recipients, patients with hematologic malignancies or those receiving immunosuppressive agents) receive a third dose of COVID-19 vaccine (**Figure 4**) (49, 89). Alongside, on 12 August, 2021, the FDA authorized an additional mRNA-based vaccine dose for certain immunocompromised individuals (90). On the other hand, the WHO notes that the benefit of a widespread use of booster vaccinations on morbidity and mortality from COVID-19 remains uncertain, also considering the alarming shortage of vaccine supplies in lower income countries (91).

## CONCLUSION

Vaccination against COVID-19 for PsC seems overall safe and effective after well-conducted vaccinations schedules (49). Yet, seroconversion rates remain lower, lagged or both across some subgroups (**Supplementary Table S2**) (19, 27, 28, 49). Although complete absence of detectable Abs after vaccination likely equates to a lack of protection, no solid data are available to establish a correlation between the protective role of vaccination and the anti-Spike Ab titer (104, 105).

Therefore, provided that PsC are comprehensively counseled about the available information on vaccine effectiveness, a tailored approach may be proposed, considering the type of malignancy

and of the specific oncologic treatment received (**Figure 4**) (89). In general, for patients with solid tumors in remission, with MPN/MDS without treatment, receiving ET, ICIs or non lymphosuppressive cytotoxic agents, no particular restrictions or time windows are advised (20, 104). Patients with lymphoid malignancies that are candidates to B cell depleting agents and those with solid tumors that are candidates to lymphosuppressive cytotoxic agents should be vaccinated prior to starting the planned regimen, if feasible. If not, vaccine doses should be planned in a time window that considers the nadir of the expected CT-induced cytopenia. Patients with hematologic malignancies already under treatment with B cell depleting agents, CD19-directed CAR-T cell therapies and HCT recipients showed very low seroconversion rates from vaccination, prompting concern that they are likely to have poor protection against COVID-19. A third vaccine dose may be proposed to such immunocompromised categories, possibly considering a time window consistent with the emerging evidence of an acceptable serological response between 9 and 12 months, after completing treatment (**Figure 4**) (20, 38, 46, 89). A similar approach may be warranted for individuals with active CLL and older patients with MM, although a case-by-case management is recommended (29, 30, 41). Findings from prospective clinical trials with a longer follow-up will further elucidate how to tailor vaccination in special populations (108-111).

A major obstacle to achieve herd immunity is vaccine hesitancy (92). In the specific population of PsC, vaccine acceptance is generally higher than in other patient populations (49, 97, 98). In an Italian report, of 914 patients eligible to the survey, only 102 refused vaccination (11.2%). The most frequently reported reasons to refuse the vaccine were concerns about vaccine-related AEs (48.1%). The identification of the reasons associated with vaccine hesitancy and refusal should be exploited to personalize educational approaches (100-103).

The reviewed data altogether suggest that PsC, especially those at the highest risk of poor seroconversion, should maintain strict preventive behaviors (e.g., FFP-2 masks), for at least 6-8 weeks after the first vaccine dose, and to not postpone the second dose, if possible (49, 110-111). Alongside, households and other close contacts of immunocompromised patients should be vaccinated (49, 89).

**Funding:** none.

**Acknowledgements:** CC and GA contributed to the literature search, conception and design of the article and drafted the first version of the manuscript. GC contributed to conception and design of the article, provided critical revisions of the manuscript and supervision. FS, JPS, JB, J-M.M and FA provided critical revisions of the manuscript. All the authors provided final approval to the submitted work.

**Competing Interests:** CC and GA have no potential conflicts of interest to disclose. FS reports consulting fees from AMGEN, Roche, Chugai, Mylan, Mundi Pharma, Leo Pharma, Pierre Fabre Oncology, Helsinn, MSD, Pfizer, BMS, all outside the submitted work. JPS has received personal fees as a speaker or consultant from Pfizer, AstraZeneca, Gilead, Leopharma, Daiichi, Mylan, BMS, MSD, Pierre Fabre, GSK (Abbvie), Roche, Novartis and Lilly. JB declares that he has no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper. As part of the Drug Development Department (DITEP), J-M.M reports: principal/sub-Investigator of Clinical Trials for Abbvie, Adaptimmune, Adlai Nortye USA Inc., Aduro Biotech, Agios Pharmaceuticals, Amgen, Argen-X Bvba, Astex Pharmaceuticals, Astra Zeneca Ab, Aveo, Basilea Pharmaceutica International Ltd., Bayer Healthcare Ag, Bbb Technologies Bv, Beigene, BicycleTx Ltd, Blueprint Medicines, Boehringer Ingelheim, Boston Pharmaceuticals, Bristol Myers Squibb, Ca, Celgene Corporation, Chugai Pharmaceutical Co, Clovis Oncology, Cullinan-Apollo, Curevac, Daiichi Sankyo, Debiopharm, Eisai, Eisai Limited, Eli Lilly, Exelixis, Faron Pharmaceuticals Ltd., Forma Therapeutics, Gamamabs, Genentech, Glaxosmithkline, H3 Biomedicine, Hoffmann La Roche Ag, Imcheck Therapeutics, Innate Pharma, Institut De Recherche Pierre Fabre, Iris Servier, Iteos Belgium SA, Janssen Cilag, Janssen Research Foundation, Kura Oncology, Kyowa Kirin Pharm. Dev, Lilly France, Loxo Oncology, Lytix Biopharma As, Medimmune, Menarini Ricerche, Merck Sharp & Dohme Chibret, Merrimack Pharmaceuticals, Merus, Millennium Pharmaceuticals, Molecular Partners Ag, Nanobiotix, Nektar Therapeutics, Novartis Pharma, Octimet Oncology Nv, Oncoethix, Oncopeptides, Orion Pharma, Ose Pharma, Pfizer, Pharma Mar,

Pierre Fabre, Medicament, Roche, Sanofi Aventis, Seattle Genetics, Sotio A.S, Syros Pharmaceuticals, Taiho Pharma, Tesaro, Turning Point Therapeutics, and Xencor; research grants from Astrazeneca, BMS, Boehringer Ingelheim, GSK, INCA, Janssen Cilag, Merck, Novartis, Pfizer, Roche, and Sanofi; and non-financial support (drug supplied) from Astrazeneca, Bayer, BMS, Boehringer Ingelheim, GSK, Medimmune, Merck, NH TherAGuiX, Pfizer, and Roche. J.-M.M. also reports other support from AstraZeneca, Bristol-Myers Squibb, Boehringer–Ingelheim, Eli Lilly Oncology, b. Hoffmann–La Roche Ltd, Merck, MSD, Pfizer, Regeneron, outside this work. Over the last 5 years, A.M. has been a Principal Investigator of Clinical Trials from the following companies: Roche/Genentech, BMS, Merck (MSD), Pfizer, Lytix pharma, Eisai, Astra Zeneca/Medimmune, Tesaro, Chugai, OSE immunotherapeutics, SOTIO, Molecular Partners, IMCheck, Pierre Fabre, and Adlai Nortye. A.M. has been a Member of Clinical Trial Steering Committees for NCT02528357 (GSK), NCT03334617 (AZ) and a Member of Data Safety and Monitoring Board: NCT02423863 (Sponsor: Oncovir), NCT03818685 (Sponsor: Centre Le´ on Be´ - rard). A.M. has been a compensated member of the following Scientific Advisory Boards: Merck Serono, eTheRNA, Lytix pharma, Kyowa Kirin Pharma, Novartis, BMS, Symphogen, Genmab, Amgen, Biothera, Nektar, Tesaro/GSK, Oncosec, Pfizer, Seattle Genetics, Astra Zeneca/Medimmune, Servier, Gritstone, Molecular Partners, Bayer, Partner Therapeutics, Sanofi, Pierre Fabre, RedX pharma, OSE Immunotherapeutics, Medicxi, HiFiBio, IMCheck, MSD, iTeos, Innate Pharma, Shattuck Labs, Medincell, Tessa Therapeutics, and Deka Biosciences, all outside the submitted work. FA reports grants from Novartis, grants from Pfizer, grants from Roche, grants from Daiichi, grants from Eli Lilly and grants from AstraZeneca, all outside the submitted work. GC served as consultant or advisor for Roche, Lilly, and Bristol-Myers Squibb, served on the speaker’s bureau for Roche, Pfizer, and Lilly, received travel funding from Pfizer and Roche, and received honoraria from Roche, Pfizer, Lilly, Novartis, AstraZeneca and SEAGEN, all outside the submitted work.

## References



- 580 1. WHO. Coronavirus disease (COVID-19) pandemic. Numbers at a glance. [Available from:  
581 <https://www.who.int/emergencies/diseases/novel-coronavirus-2019>]. Accessed on 2 September  
582 2021.
- 583 2. Sterne JAC, Murthy S, Diaz JV, Slutsky AS, Villar J, Angus DC, et al. Association Between  
584 Administration of Systemic Corticosteroids and Mortality Among Critically Ill Patients With COVID-  
585 19: A Meta-analysis. JAMA. 2020;324(13):1330-41.
- 586 3. Horby WP, Pessoa-Amorim, G, Peto L, Brightling, C.E, et al. Tocilizumab in patients admitted  
587 to hospital with COVID-19 (RECOVERY): preliminary results of a randomised, controlled, open-  
588 label, platform trial. Preprint, MedRxiv, 2021.
- 589 4. Marconi VC, Ramanan, A.V, de Bono S, Kartman, C.E, et al. Baricitinib plus Standard of Care  
590 for Hospitalized Adults with COVID-19. Preprint, MedRxiv, 2021.
- 591 5. Pan H, Peto R, Henao-Restrepo AM, Preziosi MP, Sathiyamoorthy V, Abdool Karim Q, et al.  
592 Repurposed Antiviral Drugs for Covid-19 - Interim WHO Solidarity Trial Results. N Engl J Med. 2020.
- 593 6. Kuderer NM, Choueiri TK, Shah DP, Shyr Y, Rubinstein SM, Rivera DR, et al. Clinical impact  
594 of COVID-19 on patients with cancer (CCC19): a cohort study. Lancet. 2020;395(10241):1907-18.
- 595 7. Williamson EJ, Walker AJ, Bhaskaran K, Bacon S, Bates C, Morton CE, et al. Factors  
596 associated with COVID-19-related death using OpenSAFELY. Nature. 2020;584(7821):430-6.
- 597 8. Elkrief A, Desilets A, Papneja N, Cvetkovic L, Groleau C, Lakehal YA, et al. High mortality  
598 among hospital-acquired COVID-19 infection in patients with cancer: A multicentre observational  
599 cohort study. Eur J Cancer. 2020;139:181-7.
- 600 9. Lièvre A, Turpin A, Ray-Coquard I, Le Malicot K, Thariat J, Ahle G, et al. Risk factors for  
601 Coronavirus Disease 2019 (COVID-19) severity and mortality among solid cancer patients and  
602 impact of the disease on anticancer treatment: A French nationwide cohort study (GCO-002  
603 CACOVID-19). Eur J Cancer. 2020;141:62-81.
- 604 10. de Joode K, Dumoulin DW, Tol J, Westgeest HM, Beerepoot LV, van den Berkmortel FWPJ,  
605 et al. Dutch Oncology COVID-19 consortium: Outcome of COVID-19 in patients with cancer in a  
606 nationwide cohort study. Eur J Cancer. 2020;141:171-84.



- 607 11. Saini KS, Tagliamento M, Lambertini M, McNally R, Romano M, Leone M, et al. Mortality in  
608 patients with cancer and coronavirus disease 2019: A systematic review and pooled analysis of 52  
609 studies. *Eur J Cancer*. 2020;139:43-50.
- 610 12. Yekedüz E, Utkan G, Ürün Y. A systematic review and meta-analysis: the effect of active  
611 cancer treatment on severity of COVID-19. *Eur J Cancer*. 2020;141:92-104.
- 612 13. Garassino MC, Whisenant JG, Huang LC, Trama A, Torri V, Agustoni F, et al. COVID-19 in  
613 patients with thoracic malignancies (TERAVOLT): first results of an international, registry-based,  
614 cohort study. *Lancet Oncol*. 2020;21(7):914-22.
- 615 14. Corti C, Curigliano G. Commentary: SARS-CoV-2 vaccines and cancer patients. *Ann Oncol*  
616 . 2021 Apr;32(4):569-571
- 617 15. Corti C, Crimini E, Tarantino P, Pravettoni G, Eggermont AMM, Delaloge S, et al. SARS-  
618 CoV-2 vaccines for cancer patients: a call to action. *Eur J Cancer*. 2021;148:316-27.
- 619 16. Karacin C, Eren T, Zeynelgil E, Imamoglu GI, Altinbas M, Karadag I, et al. Immunogenicity  
620 and safety of the CoronaVac vaccine in patients with cancer receiving active systemic therapy.  
621 *Future Oncol*. 2021.
- 622 17. Guglielmelli P, Mazzoni A, Maggi L, Kiros ST, Zammarchi L, Pileri S, et al. Impaired  
623 response to first SARS-CoV-2 dose vaccination in myeloproliferative neoplasm patients receiving  
624 ruxolitinib. *Am J Hematol*. 2021.
- 625 18. Terpos E, Zagouri F, Liontos M, Sklirou AD, Koutsoukos K, Markellos C, et al. Low titers of  
626 SARS-CoV-2 neutralizing antibodies after first vaccination dose in cancer patients receiving  
627 checkpoint inhibitors. *J Hematol Oncol*. 2021;14(1):86.
- 628 19. Barrière J, Chamoirey E, Adjoutah Z, Castelnau O, Mahamat A, Marco S, et al. Impaired  
629 immunogenicity of BNT162b2 anti-SARS-CoV-2 vaccine in patients treated for solid tumors. *Ann*  
630 *Oncol*. 2021;32(8):1053-5.
- 631 20. Harrington P, de Lavallade H, Doores KJ, O'Reilly A, Seow J, Graham C, et al. Single dose  
632 of BNT162b2 mRNA vaccine against SARS-CoV-2 induces high frequency of neutralising antibody  
633 and polyfunctional T-cell responses in patients with myeloproliferative neoplasms. *Leukemia*. 2021.

- 634 21. Thakkar A, Gonzalez-Lugo JD, Goradia N, Gali R, Shapiro LC, Pradhan K, et al.  
635 Seroconversion rates following COVID-19 vaccination among patients with cancer. *Cancer Cell*.  
636 2021;39(8):1081-90.e2.
- 637 22. Yatim N, Boussier J, Tetu P, Smith N, Bruel T, Charbit B, et al. Immune checkpoint inhibitors  
638 increase T cell immunity during SARS-CoV-2 infection. *Sci Adv*. 2021;7(34).
- 639 23. Iacono D, Cerbone L, Palombi L, Cavalieri E, Sperduti I, Cocchiara RA, et al. Serological  
640 response to COVID-19 vaccination in patients with cancer older than 80 years. *J Geriatr Oncol*. 2021.
- 641 24. Maneikis K, Šablauskas K, Ringelevičiūtė U, Vaitekėnaitė V, Čekauskienė R, Kryžauskaitė  
642 L, et al. Immunogenicity of the BNT162b2 COVID-19 mRNA vaccine and early clinical outcomes in  
643 patients with haematological malignancies in Lithuania: a national prospective cohort study. *Lancet*  
644 *Haematol*. 2021;8(8):e583-e92.
- 645 25. Monin L, Laing AG, Muñoz-Ruiz M, McKenzie DR, Del Molino Del Barrio I, Alaguthurai T, et  
646 al. Safety and immunogenicity of one versus two doses of the COVID-19 vaccine BNT162b2 for  
647 patients with cancer: interim analysis of a prospective observational study. *Lancet Oncol*.  
648 2021;22(6):765-78.
- 649 26. Shepherd S, Fendler A, Au L, Byrne F, Wilkinson K, Wu M, et al. 1557O - Adaptive  
650 immunity to SARS-CoV-2 infection and vaccination in cancer patients: The CAPTURE study. *Annals*  
651 *of Oncology*. 2021;32:S1129-S63.
- 652 27. Goshen-Lago T, Waldhorn I, Holland R, Szwarcwort-Cohen M, Reiner-Benaim A, Shachor-  
653 Meyouhas Y, et al. Serologic Status and Toxic Effects of the SARS-CoV-2 BNT162b2 Vaccine in  
654 Patients Undergoing Treatment for Cancer. *JAMA Oncol*. 2021.
- 655 28. Palich R, Veyri M, Marot S, Vozy A, Gligorov J, Maingon P, et al. Weak immunogenicity after  
656 a single dose of SARS-CoV-2 mRNA vaccine in treated cancer patients. *Ann Oncol*.  
657 2021;32(8):1051-3.
- 658 29. Stampfer SD, Goldwater MS, Jew S, Bujarski S, Regidor B, Daniely D, et al. Response to  
659 mRNA vaccination for COVID-19 among patients with multiple myeloma. *Leukemia*. 2021.

- 660 30. Terpos E, Trougakos IP, Gavriatopoulou M, Papassotiriou I, Sklirou AD, Ntanasis-  
661 Stathopoulos I, et al. Low neutralizing antibody responses against SARS-CoV-2 in older patients  
662 with myeloma after the first BNT162b2 vaccine dose. *Blood*. 2021;137(26):3674-6.
- 663 31. Ghandili S, Schönlein M, Lütgehetmann M, Schulze Zur Wiesch J, Becher H, Bokemeyer C,  
664 et al. Post-Vaccination Anti-SARS-CoV-2-Antibody Response in Patients with Multiple Myeloma  
665 Correlates with Low CD19+ B-Lymphocyte Count and Anti-CD38 Treatment. *Cancers (Basel)*.  
666 2021;13(15).
- 667 32. Ramasamy K, Sadler R, Jeans S, Varghese S, Turner A, Larham J, et al. COVID symptoms,  
668 testing, shielding impact on patient-reported outcomes and early vaccine responses in individuals  
669 with multiple myeloma. *Br J Haematol*. 2021.
- 670 33. Van Oekelen O, Gleason CR, Agte S, Srivastava K, Beach KF, Aleman A, et al. Highly  
671 variable SARS-CoV-2 spike antibody responses to two doses of COVID-19 RNA vaccination in  
672 patients with multiple myeloma. *Cancer Cell*. 2021;39(8):1028-30.
- 673 34. Re D, Barrière J, Chamoirey E, Delforge M, Gastaud L, Petit E, et al. Low rate of  
674 seroconversion after mRNA anti-SARS-CoV-2 vaccination in patients with hematological  
675 malignancies. *Leuk Lymphoma*. 2021;1-3.
- 676 35. Chowdhury O, Bruguier H, Mallett G, Sousos N, Crozier K, Allman C, et al. Impaired antibody  
677 response to COVID-19 vaccination in patients with chronic myeloid neoplasms. *Br J Haematol*. 2021.
- 678 36. Pimpinelli F, Marchesi F, Piaggio G, Giannarelli D, Papa E, Falcucci P, et al. Fifth-week  
679 immunogenicity and safety of anti-SARS-CoV-2 BNT162b2 vaccine in patients with multiple  
680 myeloma and myeloproliferative malignancies on active treatment: preliminary data from a single  
681 institution. *J Hematol Oncol*. 2021;14(1):81.
- 682 37. Perry C, Luttwak E, Balaban R, Shefer G, Morales MM, Aharon A, et al. Efficacy of the  
683 BNT162b2 mRNA COVID-19 vaccine in patients with B-cell non-Hodgkin lymphoma. *Blood Adv*.  
684 2021;5(16):3053-61.
- 685 38. Lim SH, Campbell N, Johnson M, Joseph-Pietras D, Collins GP, O'Callaghan A, et al.  
686 Antibody responses after SARS-CoV-2 vaccination in patients with lymphoma. *Lancet Haematol*.  
687 2021;8(8):e542-e4.

- 688 39. Herishanu Y, Avivi I, Aharon A, Shefer G, Levi S, Bronstein Y, et al. Efficacy of the BNT162b2  
689 mRNA COVID-19 vaccine in patients with chronic lymphocytic leukemia. *Blood*. 2021;137(23):3165-  
690 73.
- 691 40. Benjamini O, Rokach L, Itchaki G, Braester A, Shvidel L, Goldschmidt N, et al. Safety and  
692 efficacy of BNT162b mRNA Covid19 Vaccine in patients with chronic lymphocytic leukemia.  
693 *Haematologica*. 2021.
- 694 41. Parry H, McIlroy G, Bruton R, Ali M, Stephens C, Damery S, et al. Antibody responses after  
695 first and second Covid-19 vaccination in patients with chronic lymphocytic leukaemia. *Blood Cancer*  
696 *J*. 2021;11(7):136.
- 697 42. Eliakim-Raz N, Massarweh A, Stemmer A, Stemmer SM. Durability of Response to SARS-  
698 CoV-2 BNT162b2 Vaccination in Patients on Active Anticancer Treatment. *JAMA Oncol*. 2021.
- 699 43. Palich R, Veyri M, Vozy A, Marot S, Gligorov J, Benderra MA, et al. High seroconversion rate  
700 but low antibody titers after two injections of BNT162b2 (Pfizer-BioNTech) vaccine in patients treated  
701 with chemotherapy for solid cancers. *Ann Oncol*. 2021.
- 702 44. van der Veldt AAM, Oosting SF, Dingemans AC, Fehrmann RSN, GeurtsvanKessel C,  
703 Jalving M, et al. COVID-19 vaccination: the VOICE for patients with cancer. *Nat Med*.  
704 2021;27(4):568-9.
- 705 45. Heudel P, Favier B, Assaad S, Zrounba P, Blay JY. Reduced SARS-CoV-2 infection and  
706 death after two doses of COVID-19 vaccines in a series of 1503 cancer patients. *Ann Oncol*. 2021.
- 707 46. Ghione P, Gu JJ, Attwood K, Torka P, Goel S, Sundaram S, et al. Impaired humoral  
708 responses to COVID-19 vaccination in patients with lymphoma receiving B-cell directed therapies.  
709 *Blood*. 2021.
- 710 47. Shroff RT, Chalasani P, Wei R, Pennington D, Quirk G, Schoenle MV, et al. Immune  
711 Responses to COVID-19 mRNA Vaccines in Patients with Solid Tumors on Active,  
712 Immunosuppressive Cancer Therapy. Preprint, MedRxiv, 2021.
- 713 48. Ehmsen S, Asmussen A, Jeppesen SS, Nilsson AC, Østerlev S, Vestergaard H, et al.  
714 Antibody and T cell immune responses following mRNA COVID-19 vaccination in patients with  
715 cancer. *Cancer Cell*. 2021;39(8):1034-6.

- 716 49. Barrière J, Re D, Peyrade F, Carles M. Current perspectives for SARS-CoV-2 vaccination  
717 efficacy improvement in patients with active treatment against cancer. *Eur J Cancer*. 2021;154:66-  
718 72.
- 719 50. Anolik JH, Friedberg JW, Zheng B, Barnard J, Owen T, Cushing E, et al. B cell reconstitution  
720 after rituximab treatment of lymphoma recapitulates B cell ontogeny. *Clin Immunol*. 2007;122(2):139-  
721 45.
- 722 51. Benda M, Mutschlechner B, Ulmer H, Grabher C, Severgnini L, Volgger A, et al. Serological  
723 SARS-CoV-2 antibody response, potential predictive markers and safety of BNT162b2 mRNA  
724 COVID-19 vaccine in haematological and oncological patients. *Br J Haematol*. 2021.
- 725 52. Re D, Seitz-Polski B, Carles M, Brglez V, Graça D, et al. Brief Communication: humoral and  
726 cellular responses after a third dose of BNT162b2 vaccine in patients with lymphoid malignancies.  
727 Preprint, Research Square - Nature Portfolio2021.
- 728 53. Herzog Tzarfati K, Gutwein O, Apel A, Rahimi-Levene N, Sadovnik M, Harel L, et al.  
729 BNT162b2 COVID-19 vaccine is significantly less effective in patients with hematologic  
730 malignancies. *Am J Hematol*. 2021.
- 731 54. Laphanuwat P, Jirawatnotai S. Immunomodulatory Roles of Cell Cycle Regulators. *Front Cell*  
732 *Dev Biol*. 2019;7:23.
- 733 55. Zagouri F, Terpos E, Fiste O, Lontos M, Briasoulis A, Katsiana I, et al. SARS-CoV-2  
734 neutralizing antibodies after first vaccination dose in breast cancer patients receiving CDK4/6  
735 inhibitors. *The Breast*. 2021.
- 736 56. Desage AL, Bouleftour W, Rivoirard R, Magne N, Collard O, Fournel P, et al. Vaccination and  
737 Immune Checkpoint Inhibitors: Does Vaccination Increase the Risk of Immune-related Adverse  
738 Events? A Systematic Review of Literature. *Am J Clin Oncol*. 2021;44(3):109-13.
- 739 57. Waissengrin B, Agbarya A, Safadi E, Padova H, Wolf I. Short-term safety of the BNT162b2  
740 mRNA COVID-19 vaccine in patients with cancer treated with immune checkpoint inhibitors. *Lancet*  
741 *Oncol*. 2021;22(5):581-3.
- 742 58. Sadarangani M, Marchant A, Kollmann TR. Immunological mechanisms of vaccine-induced  
743 protection against COVID-19 in humans. *Nat Rev Immunol*. 2021;21(8):475-84.

59. Mairhofer M, Kausche L, Kaltenbrunner S, Ghanem R, Stegemann M, Klein K, et al. Humoral and cellular immune responses in SARS-CoV-2 mRNA-vaccinated patients with cancer. *Cancer Cell*. 2021.
60. Lipsitch M, Swerdlow DL, Finelli L. Defining the Epidemiology of Covid-19 - Studies Needed. *N Engl J Med*. 2020;382(13):1194-6.
61. Sette A, Crotty S. Adaptive immunity to SARS-CoV-2 and COVID-19. *Cell*. 2021;184(4):861-80.
62. Rydyznski Moderbacher C, Ramirez SI, Dan JM, Grifoni A, Hastie KM, Weiskopf D, et al. Antigen-Specific Adaptive Immunity to SARS-CoV-2 in Acute COVID-19 and Associations with Age and Disease Severity. *Cell*. 2020;183(4):996-1012.e19.
63. Tan AT, Linster M, Tan CW, Le Bert N, Chia WN, Kunasegaran K, et al. Early induction of functional SARS-CoV-2-specific T cells associates with rapid viral clearance and mild disease in COVID-19 patients. *Cell Rep*. 2021;34(6):108728.
64. Dan JM, Mateus J, Kato Y, Hastie KM, Yu ED, Faliti CE, et al. Immunological memory to SARS-CoV-2 Accessed for up to 8 months after infection. *Science*. 2021;371(6529).
65. Krammer F. A correlate of protection for SARS-CoV-2 vaccines is urgently needed. *Nat Med*. 2021;27(7):1147-8.
66. Nanduri S, Pilishvili T, Derado G, Soe M, Dollard P, Wu H, et al. Effectiveness of Pfizer-BioNTech and Moderna Vaccines in Preventing SARS-CoV-2 Infection Among Nursing Home Residents Before and During Widespread Circulation of the SARS-CoV-2 B.1.617.2 (Delta) Variant 2021 [Available from: <https://www.cdc.gov/mmwr/volumes/70/wr/mm7034e3.htm>]. Accessed on 2 September 2021.
67. Rosenberg ES, Holtgrave DR, Dorabawila V, Conroy M, Greene D, Lutterloh E, et al. The Centers for Disease Control and Prevention (CDC): Morbidity and Mortality Weekly Report (MMWR) - New COVID-19 Cases and Hospitalizations Among Adults, by Vaccination Status 2021 [Available from: <https://www.cdc.gov/mmwr/volumes/70/wr/mm7034e1.htm>]. Accessed on 2 September 2021.
68. Tenforde MW, Self WH, Naioti EA, Ginde AA, Douin DJ, Olson SM, et al. Centers For Disease Control and Prevention (CDC): Morbidity and Mortality Weekly Report (MMWR) - Sustained

- Effectiveness of Pfizer-BioNTech and Moderna Vaccines Against COVID-19 Associated Hospitalizations Among Adults 2021 [Available from: <https://stacks.cdc.gov/view/cdc/108947>]. Accessed on 2 September 2021.
69. Garassino MC, Giesen N, Grivas P, Jordan K, Lucibello F, Mir O, et al. ESMO Statements for Vaccination against COVID-19 in patients with cancer. [Available from: <https://bit.ly/3swNx9G> 2020]. Accessed on 2 September 2021.
70. Bilich T, Roerden M, Maringer Y, Nelde A, Heitmann JS, Dubbelaar ML, et al. Preexisting and Post-COVID-19 Immune Responses to SARS-CoV-2 in Patients with Cancer. *Cancer Discov.* 2021;11(8):1982-95.
71. Polack FP, Thomas SJ, Kitchin N, Absalon J, Gurtman A, Lockhart S, et al. Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine. *N Engl J Med.* 2020;383(27):2603-15.
72. Gounant V, Ferré VM, Soussi G, Charpentier C, Flament H, Fidouh N, et al. Efficacy of SARS-CoV-2 vaccine in thoracic cancer patients: a prospective study supporting a third dose in patients with minimal serologic response after two vaccine doses. Preprint, MedRxiv2021.
73. Mansi L, Spehner L, Daguindau E, Bouiller K, Almotlak H, Stein U, et al. Study of the SARS-CoV-2-specific immune T-cell responses in COVID-19-positive cancer patients. *Eur J Cancer.* 2021;150:1-9.
74. Liu X, Shaw RH, Stuart ASV, Greenland M, Aley PK, Andrews NJ, et al. Safety and immunogenicity of heterologous versus homologous prime-boost schedules with an adenoviral vectored and mRNA COVID-19 vaccine (Com-COV): a single-blind, randomised, non-inferiority trial. *Lancet.* 2021.
75. Potsch DV, Camacho LA, Tuboi S, Villar LM, Miguel JC, Ginuino C, et al. Vaccination against hepatitis B with 4-double doses increases response rates and antibodies titers in HIV-infected adults. *Vaccine.* 2012;30(41):5973-7.
76. Branagan AR, Duffy E, Gan G, Li F, Foster C, Verma R, et al. Tandem high-dose influenza vaccination is associated with more durable serologic immunity in patients with plasma cell dyscrasias. *Blood Adv.* 2021;5(5):1535-9.



- 799 77. Sodhi JS, Raja W, Zargar SA, Showkat A, Parveen S, Nisar S, et al. The efficacy of  
800 accelerated, multiple, double-dose hepatitis B vaccine against hepatitis B virus infection in cancer  
801 patients receiving chemotherapy. *Indian J Gastroenterol*. 2015;34(5):372-9.
- 802 78. Mulligan MJ, Lyke KE, Kitchin N, Absalon J, Gurtman A, Lockhart S, et al. Phase I/II study of  
803 COVID-19 RNA vaccine BNT162b1 in adults. *Nature*. 2020;586(7830):589-93.
- 804 79. Walsh EE, Frenck RW, Falsey AR, Kitchin N, Absalon J, Gurtman A, et al. Safety and  
805 Immunogenicity of Two RNA-Based Covid-19 Vaccine Candidates. *N Engl J Med*. 2020.
- 806 80. GOV.UK: OCTAVE DUO - New study to test third COVID-19 vaccine for people with  
807 weakened immune systems [press release, available at: <https://bit.ly/3zTQpQm>], 2021. Accessed  
808 on 2 September 2021.
- 809 81. Kearns P, Siebert S, Willicombe, M, Gaskell C, Kirkham A, et al. Examining the  
810 immunological effects of COVID-19 vaccination in patients with conditions potentially leading to  
811 diminished immune response capacity – the OCTAVE Trial. Preprints, THE LANCET, 24 August  
812 2021.
- 813 82. Karaba AH, Zhu X, Liang T, Wang KH, Rittenhouse, A.G, et al. A Third Dose of SARS-CoV-  
814 2 Vaccine Increases Neutralizing Antibodies Against Variants of Concern in Solid Organ Transplant  
815 Recipients. Preprint, MedRxiv, 14 August 2021.
- 816 83. Bar-On YM, Goldberg Y, Mandel M, Bodenheimer O, Freedman L, Kalkstein N, et al.  
817 Protection of BNT162b2 Vaccine Booster against Covid-19 in Israel. *N Engl J Med*. 2021.
- 818 84. Longlune N, Nogier MB, Miedougé M, Gabilan C, Cartou C, Seigneure B, et al. High  
819 immunogenicity of a messenger RNA based vaccine against SARS-CoV-2 in chronic dialysis  
820 patients. *Nephrol Dial Transplant*. 2021.
- 821 85. Kamar N, Abravanel F, Marion O, Couat C, Izopet J, Del Bello A. Three Doses of an mRNA  
822 Covid-19 Vaccine in Solid-Organ Transplant Recipients. *N Engl J Med*. 2021;385(7):661-2.
- 823 86. Werbel WA, Boyarsky BJ, Ou MT, Massie AB, Tobian AAR, Garonzik-Wang JM, et al. Safety  
824 and Immunogenicity of a Third Dose of SARS-CoV-2 Vaccine in Solid Organ Transplant Recipients:  
825 A Case Series. *Ann Intern Med*. 2021.



- 826 87. Hall VG, Ferreira VH, Ku T, Ierullo M, Majchrzak-Kita B, Chaparro C, et al. Randomized Trial  
827 of a Third Dose of mRNA-1273 Vaccine in Transplant Recipients. *N Engl J Med*. 2021.
- 828 88. Benotmane I, Gautier G, Perrin P, Olagne J, Cognard N, Fafi-Kremer S, et al. Antibody  
829 Response After a Third Dose of the mRNA-1273 SARS-CoV-2 Vaccine in Kidney Transplant  
830 Recipients With Minimal Serologic Response to 2 Doses. *JAMA*. 2021.
- 831 89. NCCN - Recommendations of the National Comprehensive Cancer Network COVID-19  
832 Vaccination Advisory Committee\* Version 4.0 08/30/2021 30 August 2021 [Available from:  
833 <https://bit.ly/3ihKQoX>]. Accessed on 2 September 2021.
- 834 90. FDA. Coronavirus (COVID-19) Update: The Food and Drug Administration Authorizes  
835 Additional Vaccine Dose for Certain Immunocompromised Individuals 2021 [Available from:  
836 <https://bit.ly/3gy5YX8>]. Accessed on 2 September 2021.
- 837 91. World Health Organization (WHO) - Interim statement on COVID-19 vaccine booster doses  
838 2021 [Available from: <https://bit.ly/3DzTpUX>]. Accessed on 2 September 2021.
- 839 92. Lawrence GL, Hull BP, MacIntyre CR, McIntyre PB. Reasons for incomplete immunisation  
840 among Australian children. A national survey of parents. *Aust Fam Physician*. 2004;33(7):568-71.
- 841 93. Lazarus JV, Ratzan SC, Palayew A, Gostin LO, Larson HJ, Rabin K, et al. A global survey of  
842 potential acceptance of a COVID-19 vaccine. *Nat Med*. 2021;27(2):225-8.
- 843 94. Stoeklé HC, Sekkate S, Angellier E, Hervé C, Beuzeboc P. Refusal of anti-coronavirus  
844 disease 2019 vaccination in cancer patients: Is there a difference between the sexes? *Eur J Cancer*.  
845 2021;155:54-5.
- 846 95. Daly M, Jones A, Robinson E. Public Trust and Willingness to Vaccinate Against COVID-19  
847 in the US From October 14, 2020, to March 29, 2021. *JAMA*. 2021;325(23):2397-9.
- 848 96. Fisher KA, Bloomstone SJ, Walder J, Crawford S, Fouayzi H, Mazor KM. Attitudes Toward a  
849 Potential SARS-CoV-2 Vaccine : A Survey of U.S. Adults. *Ann Intern Med*. 2020;173(12):964-73.
- 850 97. Chun JY, Kim SI, Park EY, Park SY, Koh SJ, Cha Y, et al. Cancer Patients' Willingness to  
851 Take COVID-19 Vaccination: A Nationwide Multicenter Survey in Korea. *Cancers (Basel)*.  
852 2021;13(15).

98. Barrière J, Gal J, Hoch B, Cassuto O, Leysalle A, Chamorey E, et al. Acceptance of SARS-CoV-2 vaccination among French patients with cancer: a cross-sectional survey. *Ann Oncol*. 2021;32(5):673-4.
99. Di Noia V, Renna D, Barberi V, Di Civita M, Riva F, Costantini G, et al. The first report on coronavirus disease 2019 (COVID-19) vaccine refusal by patients with solid cancer in Italy: Early data from a single-institute survey. *Eur J Cancer*. 2021;153:260-4.
100. Opel DJ, Henrikson N, Lepere K, Hawkes R, Zhou C, Dunn J, et al. Previsit Screening for Parental Vaccine Hesitancy: A Cluster Randomized Trial. *Pediatrics*. 2019;144(5).
101. Curigliano G, Eggermont AMM. Adherence to COVID-19 vaccines in cancer patients: promote it and make it happen! *Eur J Cancer*. 2021;153:257-9.
102. Breza E, Stanford FC, Alsan M, Alsan B, Banerjee A, Chandrasekhar AG, et al. Effects of a large-scale social media advertising campaign on holiday travel and COVID-19 infections: a cluster randomized controlled trial. *Nat Med*. 2021.
103. Spano JP, Barre-Sinoussi F, Kieny MP, Marcelin AG, Blay JY. [COVID-19 vaccination for cancer patients: Medical and ethical need]. *Bull Cancer*. 2021;108(3):225-7.
104. Griffiths EA, Segal BH. Immune responses to COVID-19 vaccines in patients with cancer: Promising results and a note of caution. *Cancer Cell*. 2021;39(8):1045-7.
105. Khoury DS, Cromer D, Reynaldi A, Schlub TE, Wheatley AK, Juno JA, et al. Neutralizing antibody levels are highly predictive of immune protection from symptomatic SARS-CoV-2 infection. *Nat Med*. 2021;27(7):1205-11.
106. Andreano E, Nicastrì E, Paciello I, Pileri P, Manganaro N, Piccini G, et al. Extremely potent human monoclonal antibodies from COVID-19 convalescent patients. *Cell*. 2021;184(7):1821-35.e16.
107. Kared H, Redd AD, Bloch EM, Bonny TS, Sumatoh H, Kairi F, et al. SARS-CoV-2-specific CD8+ T cell responses in convalescent COVID-19 individuals. *J Clin Invest*. 2021;131(5).
108. Loubet P, Wittkop L, Tartour E, Parfait B, Barrou B, Blay JY, et al. A French cohort for assessing COVID-19 vaccine responses in specific populations. *Nat Med*. 2021;27(8):1319-21.

- 880 109. Desai A, Gainor JF, Hegde A, Schram AM, Curigliano G, Pal S, et al. COVID-19 vaccine  
 881 guidance for patients with cancer participating in oncology clinical trials. *Nat Rev Clin Oncol.*  
 882 2021;18(5):313-9.
- 883 110. Di Noia V, Pimpinelli F, Renna D, Barberi V, Maccallini MT, Gariazzo L, et al. Immunogenicity  
 884 and safety of COVID-19 vaccine BNT162b2 for patients with solid cancer: a large cohort prospective  
 885 study from a single institution. *Clin Cancer Res.* 2021.
- 886 111. Mair M, Berger J, Berghoff A, Starzer A, Ortmayr G, Pühr H, et al. Humoral Immune Response  
 887 in Hematooncological Patients and Health Care Workers Who Received SARS-CoV-2 Vaccinations.  
 888 *JAMA Oncol.* 2021.

889

## 890 **Figure Captions**

891

892 **Figure 1. Investigated strategies against SARS-CoV-2.** Main mechanisms of viral entry into host  
 893 cells are depicted, alongside anti-SARS-CoV-2 passive and active immune strategies. Strategy 1  
 894 and 2: soluble RBD mimetics or anti-ACE2 scFvs may hide ACE2 receptors from spike proteins,  
 895 preventing viral entry. RBD targeting may be achieved via either monoclonal Ab (i.e., casirivimab-  
 896 imdevimab) or vaccine-induced Ab. In addition, vaccination also promotes the emergence of cellular  
 897 anti-SARS-CoV2 adaptive immune responses, leading to killing of viral-infected cells. Abbreviations:  
 898 SARS-CoV-2, Severe Acute Respiratory Syndrome Coronavirus-2; RBD, Receptor Binding Domain;  
 899 ACE2, Angiotensin-converting enzyme 2; mRNA, messenger ribonucleic acid; S protein, spike  
 900 protein; Abs, antibodies; scFv, single chain variable fragment. Created with biorender.com.

901

902 **Figure 2. PRISMA flow diagram of the study.** Abbreviations: pts, patients; SARS-CoV-2, severe  
 903 acute respiratory syndrome coronavirus 2; n, number.

904

905 **Figure 3. Estimated spectrum of COVID-19 vaccine efficacy for patients with cancer,**  
 906 **according to cancer types and therapies.** Specific patient populations, especially those with  
 907 hematologic malignancies receiving B-cell targeted agents, stem cell transplantation or CAR-T cell


treatment, may not mount a protective response. Further research is warranted to clarify the vaccine-induced immune response in a number of cancer types and regimen, particularly in those receiving targeted therapy and investigational drugs. # B-cell targeted agents include anti-CD20 agents (e.g., rituximab), anti-CD38 therapy, BCMA targeted agents, and Bruton tyrosine kinase inhibitors. Abbreviations: JAKi, janus kinase inhibitor; ET, endocrine therapy; ICI, immune checkpoint inhibitors; tx, therapies; MPN, myeloproliferative neoplasms; MDS, myelodysplastic syndromes; CLL, chronic lymphocytic leukemia; CT, chemotherapy; CAR, chimeric antigen receptor; mo, months; pts, patients; MM, multiple myeloma; BCMA, B-cell maturation antigen. Created with biorender.com.

**Figure 4. Consensus recommendations on patients with cancer who should be prioritized for a third dose.** Within each box, the subgroups which should receive a third dose with priority, as they may represent the most immunocompromised individuals per each macro category (**bold**), are reported. The CDC recommends the additional dose of an mRNA COVID-19 vaccine be administered at least four weeks after a second dose of the Pfizer-BioNTech or Moderna vaccine. For people who received the Pfizer-BioNTech or Moderna COVID-19 vaccine series, a third dose of the same mRNA vaccine should be used if possible. If the same mRNA vaccine isn't available for the third dose administration or is unknown, either mRNA COVID-19 vaccine may be used. The use of antibody titers to determine if patients should receive the third dose is not recommended (outside of a research study). Abbreviations: pts, patients; SCT, stem cell transplant; GvHD, Graft-versus-host disease; HIV, Human Immunodeficiency Virus; CDC, Centers for Disease Control and Prevention. Source: National Comprehensive Cancer Network (NCCN) COVID-19 Vaccination Advisory Committee. Version 4.0 08/30/2021. Created with biorender.com.

## PASSIVE IMMUNITY


## ACTIVE IMMUNITY

### Viral entry mechanism of SARS-CoV-2




### Targeting ACE2 receptors

**Strategy 1**  
Soluble RBD mimetic




**Strategy 2**  
scFv against ACE2




### Targeting RBD of Spike protein

i.e.,  
**casirivimab - imdevimab**



### Vaccine





Identification

Screening

Eligibility

Inclusion





## Patients with cancer who should be prioritized for a third dose

### Hematopoietic cell transplant and CAR-T cell therapy

- Pts who are  $\leq 2$  years post-procedure
- Allogeneic SCT recipients receiving immunosuppressive therapy or with history of GvHD, regardless of the time post-transplant

### All hematologic malignancies

### Solid tumor malignancies

- Pts who received cancer therapy within 1-year of the initial vaccine administration
- Pts newly diagnosed with cancer or recurrent cancer who will receive cancer therapy

### Other

- Pts with cancer AND other concurrent immunocompromising conditions (e.g. HIV or autoimmune diseases).
- Pts treated with systemic corticosteroids or other concomitant immunosuppressive agents